arXiv:1605.06230v1 [math.AG] 20 May 2016

PROJECTIONS OF VERONESE SURFACE AND MORPHISMS FROM PROJECTIVE PLANE TO GRASSMANNIAN

A. EL MAZOUNI, F. LAYTIMI, AND D.S. NAGARAJ

Abstract. In this note we describe the image of $\mathbb{P}^2$ in $Gr(2,\mathbb{C}^4)$ under a morphism given by a rank two vector bundle on $\mathbb{P}^2$ with Chern classes $(2,2)$.

Keywords: Projective plane; Vector bundles; Morphisms; Grassmannian; Veronese surface.

1. Introduction

We denote by $\mathbb{P}^2$ the projective plane over the field $\mathbb{C}$ of complex numbers and by $Gr(2,\mathbb{C}^4)$ the Grassmannian variety of two dimensional quotient spaces of $\mathbb{C}^4$.

Let Q be a rank two vector bundle on $\mathbb{P}^2$ generated by global sections. Then Q can be generated by at most four linearly independent section. Assume that Q is generated by four linearly independent sections but is not generated by less number of sections. If we fix a set S of four linearly independent global sections generating Q we get a non constant morphism

$$\phi_S : \mathbb{P}^2 \to Gr(2,\mathbb{C}^4).$$

The aim of this article is to study the properties of the image of $\mathbb{P}^2$ under such a morphism ϕ_S.

According to a result of Tango [11], if ϕ_S is an embedding then the pair of Chern classes $(c_1(Q), c_2(Q))$ is either $(2,1)$ or $((2,3)$.

It is interesting to know what is the image of $\mathbb{P}^2$ under a general morphism $\phi_S : \mathbb{P}^2 \to Gr(2,\mathbb{C}^4)$. For example, What are the possible singularities of the image ? What are the defining equations of the image?

In this article we focus on the image of $\mathbb{P}^2$ in $Gr(2,\mathbb{C}^4)$ associated to a globally generated vector bundle Q with Chern class pair $(c_1(Q), c_2(Q)) = (2,2)$. More precisely we have the following Theorem(5.1):

1991 *Mathematics Subject Classification.* 14F17.

Theorem 1.1. *Let $\phi : \mathbb{P}^2 \to Gr(2, \mathbb{C}^4)$ be a morphism. Assume that $c_1(Q) = 2$ and $c_2(Q) = 2$, where Q is the pull back by ϕ of the universal rank two quotient bundle on $Gr(2, \mathbb{C}^4)$. Then the image of $\mathbb{P}^2$ in $Gr(2, \mathbb{C}^4)$ is*

a) *either a complete intersection of two independent hyperplanes,*
b) *or is a complete intersection of a hyperplane and a quadric.*

Here a hyperplane(respectively, quadric) means a divisor in the class of the ample generator (respectively, twice the class of the ample generator) of the the Picard group of $Gr(2, \mathbb{C}^4)$. In case of a) the image is isomorphic to a cone in $\mathbb{P}^3$ over a conic. In the case of b) the image is isomorphic to base locus of a pencil consisting of singular quadrics in $\mathbb{P}^4$ of rank 3 and 4. More over the image surface is singular exactly along a line of $\mathbb{P}^4$ contained in the surface.

2. Special Projections of Veronese Surface

Our study of the image of $\mathbb{P}^2$ in $Gr(2, \mathbb{C}^4)$ under the morphism given by a rank two vector bundle on $\mathbb{P}^2$ with Chern classes $(2, 2)$, depends on the study of the projections of Veronese surface V in $\mathbb{P}^5$ from a special point not on the surface or a special line not intersecting the Veronese surface. Here special point means a point on the secant variety of V and a special line means a line contained in the secant variety. In this section we recall some facts about the Veronese surface (see, [2], [10] and [6] for details) and deduce some results about special projections.

Veronese surface is the only non-degenerate (i.e., not contained in a hyperplane) non-singular surface in $\mathbb{P}^5$ which can be projected isomorphically to $\mathbb{P}^4$. A general projection has this property. It is well known that the Veronese surface is the unique closed orbit for the natural action of the algebraic group $PGL(3, \mathbb{C}^3)$ on $\mathbb{P}^5$. In fact if we identify $\mathbb{P}^5$ with the space of conics in $\mathbb{P}^2$, there are three orbits namely,

- the set of all non-singular quadrics
- the set of all pairs of distinct lines
- the set of all double lines.

The set of double lines is the Veronese surface. (See, [10, p.120-21]).

The following remarks shed light on the image of Veronese surface under special projections.

Remark 2.1. *Let V be a Veronese surface in $P(H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(2)))$ and $sec(V)$ be its secant variety. Let $p \in sec(V) \setminus V$ be a point. Then the the projection of $\mathbb{P}^5 \setminus \{p\}$ from p to a hyperplane $H \simeq \mathbb{P}^4$ in $\mathbb{P}^5$ not containing p maps V onto a singular surface V_p of degree four. V_p is cut out by intersection of quadrics of a linear pencil of quadrics in $\mathbb{P}^4$, and*

each qudaric in the pencil is either has rank three or four. More over V_p is singular along a line L_p of $\mathbb{P}^4$. The projection map $f : V \to V_p$ induces an isomorphism from $V \setminus f^{-1}(L_p)$ to $V_p \setminus L_p$, and $f^{-1}(L_p)$ is a conic and the map f restricted to $f^{-1}(L_p)$ to L_p is a ramified covering of degree two.

See [2, p. 366, 10.5.5.] for details. The only thing missing there is the statement that V_p is cut out by intersection of quadrics of $\mathbb{P}^4$, in a linear pencil of quadrics in $\mathbb{P}^4$, and each quadric in the pencil is either has rank three or four. This can be seen in several ways. One of the way is to use the fact that the set of all pairs of distinct lines is one orbit for the action for $PGL(3, \mathbb{C}^3)$ namely $sec(V) \setminus V$ and to prove that required property holds for the projection *corresponds to a particular point* $p \in sec(V) \setminus V$. The image morphism $\mathbb{P}^2 \to \mathbb{P}^4$ given by

$$(x, y, z) \mapsto (z^2, -xy, -y^2 + yz, x^2, xy - xz)$$

is one such projection. The image is cut out by the pencil of quadrics $\lambda(Z_1Z_4 - Z_2Z_3) + \mu((Z_1 + Z_4)^2 - Z_0Z_3)$, where Z_i $(0 \leq i \leq 4)$ are the homogeneous coordinate functions on $\mathbb{P}^4$.

Remark 2.2. *Let V be a Veronese surface in $P(H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(2) \simeq \mathbb{P}^5$ and $sec(V)$ be its secant variety. Let $\ell \subset sec(V)$ be a line of $\mathbb{P}^5$ such that $V \cap \ell = \emptyset$. Then the the projection of $\mathbb{P}^5 \setminus \ell$ from ℓ to a linear subspace $L \simeq \mathbb{P}^3$ in $\mathbb{P}^5$ not meeting ℓ maps V onto a qudaric surface V_ℓ, which is a cone over a quadric. The projection map $f : V \to V_\ell$ is generically two to one.*

Existence of lines in $sec(V)$ with the above property follows from [2, p. 361, 10.4.]. For example, for a fixed line $L_0 \subset \mathbb{P}^2$ the plane $[L_0](\subset sec(V)) \subset \mathbb{P}^5$ defined by

$$[L_0] := \{L_0 \cup M | M \subset \mathbb{P}^2, \text{a line}\}$$

meet the Veronese surface V in a single point l_0. Here we have identified V as set of all lines in $\mathbb{P}^2$ under the map $L \mapsto L^2$. Under this identification $sec(V)$ corresponds to set of reducible conics. Hence any line in the plane $[L_0]$ not passing through $l_0$has the required property.

3. Morphisms from $\mathbb{P}^2$ to $Gr(2, \mathbb{C}^4)$.

For a vector bundle E on $\mathbb{P}^2$ the bundle $E \otimes \mathcal{O}_{\mathbb{P}^2}(k)$ is denoted by $E(k)$.

A globally generated rank two vector bundle Q on $\mathbb{P}^2$ can be generated by at most four linearly independent global sections. If we take a

set of generators consisting of at most four sections of Q we get a morphism from $\mathbb{P}^2$ to $Gr(2,\mathbb{C}^4)$. Note that the bundle Q is generated by two linearly independent sections if and only if $Q \simeq \mathcal{O}^2_{\mathbb{P}^2}$. This happens if and only the morphism from $\mathbb{P}^2$ to $Gr(2,\mathbb{C}^4)$ is constant. Moreover the bundle Q generated by three linearly independent sections if and only if the morphism from $\mathbb{P}^2$ to $Gr(2,\mathbb{C}^4)$ factors through a linear $\mathbb{P}^2 \simeq Gr(2,\mathbb{C}^3)$ contained in $Gr(2,\mathbb{C}^4)$.

Definition 3.1. Let Q be a rank two vector bundle on $\mathbb{P}^2$ generated by global sections. Assume that Q cannot be generated by less than four independent sections. If S is a set of four independent global sections generating Q, then we get morphism

$$\phi_S : \mathbb{P}^2 \to Gr(2,\mathbb{C}^4).$$

We call such a morphism ϕ_S a non special morphism. (Generally, we use the the notation ϕ instead of ϕ_S.)

Remark 3.2. *1) Let $\phi_S : \mathbb{P}^2 \to Gr(2,\mathbb{C}^4)$ be a non special morphism obtained from a rank two vector bundle Q. Then the pull back to $\mathbb{P}^2$ by ϕ_S of the universal quotient bundle on $Gr(2,\mathbb{C}^4)$ is equal to Q. Since the morphism ϕ_S is non special $\det(Q) = \mathcal{O}_{\mathbb{P}^2}(d)$ for some $d > 0$, i.e., $c_1(Q) > 0$. As Q is generated by sections we see that $c_2(Q) \leq 0$. A rank two bundle Q generated by sections has $c_2(Q) = 0$ implies $Q = \mathcal{O}_{\mathbb{P}^2} \oplus \mathcal{O}_{\mathbb{P}^2}(d)$. This implies $\phi_S : \mathbb{P}^2 \to Gr(2,\mathbb{C}^4)$ is a special morphism, a contradiction to the assumption. Thus we must have $c_2(Q) > 0$.*

2) Let $\phi_S : \mathbb{P}^2 \to Gr(2,\mathbb{C}^4)$ be a non special morphism as above. If $p : Gr(2,\mathbb{C}^4) \to \mathbb{P}^5$ is the Plucker imbedding, then note that the morphism

$$p \circ \phi_S : \mathbb{P}^2 \to \mathbb{P}^5$$

may not be non degenerate in the usual sense. In other words the image of $\mathbb{P}^2$ in $\mathbb{P}^5$ under $p \circ \phi_S$ may be very well contained in a hyperplane of $\mathbb{P}^5$.

For the study of non special morphisms one need to know what are the globally generated rank two vector bundles on $\mathbb{P}^2$. In our previous paper [1] we obtained some partial results about the possible Chern classes $(c_1(Q), c_2(Q))$ of rank two vector bundles Q on $\mathbb{P}^2$ generated by four sections. In [5] Ph.Ellia determined the Chern classes of rank two globally generated vector bundles on $\mathbb{P}^2$. His result gives the complete numerical characterization of such bundles. Note that a rank two globally generated vector bundles on $\mathbb{P}^2$ can be generated by 4 sections and hence gives rise to a morphism from $\mathbb{P}^2$ to $Gr(2,\mathbb{C}^4)$. Globally generated vector bundle on projective spaces with special Chern classes are studied in [3] and [4].

According to a result of Tango [11], if ϕ_S is a non special imbedding then the Chern class pair $(c_1(Q), c_2(Q))$, of Q is either $(2,1)$ or $(2,3)$.

The aim of this note is to investigate the properties of the image of $\mathbb{P}^2$ under ϕ_S for the case $c_1(Q) = 2$, $c_2(Q) = 2$. For such a bundle we have the following:

Lemma 3.3. *Let Q be a rank two vector bundle on $\mathbb{P}^2$ with $c_1(Q) = 2$, $c_2(Q) = 2$. If Q is generated by sections then Q is semi-stable.*

Proof: Assume that Q is not semi-stable, then by Lemma (3.1) of [8] $h^0(Q(-2)) \neq 0$. Let k be the largest integer such that $h^0(Q(-k)) \neq 0$. Note that $k \geq 2$ and there is an exact sequence of sheaves

$$0 \to \mathcal{O}_{\mathbb{P}^2} \to Q(-k) \to I_Z(-2k+2) \to 0, \tag{1}$$

where I_Z is the ideal sheaf of a zero-dimensional closed sub-scheme Z of length $k^2 - 2k + 2 = (k-1)^2 + 1$. Now tensoring the exact sequence (1) with the line bundle $\mathcal{O}_{\mathbb{P}^2}(k)$ we get the following exact sequence

$$0 \to \mathcal{O}_{\mathbb{P}^2}(k) \to Q \to I_Z(-k+2) \to .0 \tag{2}$$

Since Q generated by sections $k = 2$ and $Z = \emptyset$. This implies $c_2(Q) = 0$, a contradiction. This contradiction proves the lemma. □

Lemma 3.4. *Let Q be a semi-stable rank two vector bundle on $\mathbb{P}^2$ with $c_1(Q) = 2$, $c_2(Q) = 2$. Then Q is generated by four independent sections and hence there is a surjective morphism of bundles*

$$\mathcal{O}^4_{\mathbb{P}^2} \to Q \to 0$$

which determines a morphism $\phi : \mathbb{P}^2 \to Gr(2, \mathbb{C}^4)$.

Proof: If Q is a rank two vector bundle on $\mathbb{P}^2$ with $c_1(Q) = 2$, $c_2(Q) = 2$ then $Q(-1)$ has $c_1(Q(-1)) = 0$, $c_2(Q(-1)) = 1$. Now Riemann-Roch theorem together with Proposition (7.1) and Theorem (7.4) of [7] we see that $h^0(Q(-1)) = 1$. Hence there is an exact sequence

$$0 \to \mathcal{O}_{\mathbb{P}^2}(1) \to Q \to I_p(1) \to 0, \tag{3}$$

where I_p is the ideal sheaf of a point $p \in \mathbb{P}^2$. From the exact sequence (3) we see that $h^0(Q) = 5$ and Q is generated by sections. Since Q is a rank two vector bundle on $\mathbb{P}^2$ generated by section we see that Q is generated by 4 independent sections. Hence we obtain a surjective morphism of bundles

$$\mathcal{O}^4_{\mathbb{P}^2} \to Q \to 0$$

which determines a morphism $\phi : \mathbb{P}^2 \to Gr(2, \mathbb{C}^4)$ as required. □

Let Q be rank two vector bundle on $\mathbb{P}^2$ with $c_1(Q) = 2$, $c_2(Q) = 2$. By Lemma (3.4) Q determines a morphism $\phi : \mathbb{P}^2 \to Gr(2, \mathbb{C}^4)$.

Lemma 3.5. *Let Q be a semi-stable rank two vector bundle on $\mathbb{P}^2$ with $c_1(Q) = 2$, $c_2(Q) = 2$. Then there is an exact sequence*

$$0 \to \mathcal{O}_{\mathbb{P}^2}(-1) \to \mathcal{O}_{\mathbb{P}^2}^2 \oplus \mathcal{O}_{\mathbb{P}^2}(1) \to Q \to 0 \tag{4}$$

of vector bundles on $\mathbb{P}^2$.

Proof: From the proof of Lemma (3.4) it follows that Q fits into an exact sequence (3) and the induced map

$$\mathrm{H}^0(\mathbb{P}^2, Q) \to \mathrm{H}^0(\mathbb{P}^2, I_p(1))$$

of cohomology groups is surjective. Observe that if $p \in \mathbb{P}^2$ then $\mathrm{H}^0(\mathbb{P}^2, I_p(1))$ is two dimensional vector space and the natural map

$$\mathrm{H}^0(\mathbb{P}^2, I_p(1)) \otimes \mathcal{O}_{\mathbb{P}^2} \to I_p(1)$$

is surjective. Thus from (3) and the above observations we see that there is a surjection of vector bundles

$$\mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}^2 \to Q \to 0.$$

A simple Chern class computation will show that the kernel of this surjection is equal to the line bundle $\mathcal{O}_{\mathbb{P}^2}(-1)$ and hence we get the existence of required exact sequence (4). □

4. Explicit constructions

If X, Y and Z is the standard basis of $\mathrm{H}^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(1))$, then the global section (X, Y, Z^2) of the vector bundle $\mathcal{O}_{\mathbb{P}^2}(1)^2 \oplus \mathcal{O}_{\mathbb{P}^2}(2)$ is nowhere vanishing on $\mathbb{P}^2$. Thus the bundle map

$$0 \to \mathcal{O}_{\mathbb{P}^2}(-1) \to \mathcal{O}_{\mathbb{P}^2}^2 \oplus \mathcal{O}_{\mathbb{P}^2}(1)$$

given by (X, Y, Z^2) is injective. If Q is the cokernel of this injection then

$$0 \to \mathcal{O}_{\mathbb{P}^2}(-1) \overset{(X,Y,Z^2)}{\longrightarrow} \mathcal{O}_{\mathbb{P}^2}^2 \oplus \mathcal{O}_{\mathbb{P}^2}(1) \to Q \to 0 \tag{5}$$

is an exact sequence of vector bundles on $\mathbb{P}^2$ with rank of Q two and $c_1(Q) = 2$, $c_2(Q) = 2$. From cohomology exact sequence associated to (5) we see that

$$\mathrm{H}^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}^2 \oplus \mathcal{O}_{\mathbb{P}^2}(1)) \simeq \mathrm{H}^0(\mathbb{P}^2, Q). \tag{6}$$

The vector space $\mathrm{H}^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}^2 \oplus \mathcal{O}_{\mathbb{P}^2}(1))$ is equal to

$$\oplus_{i=1}^5 \mathbb{C} v_i,$$

where $v_1 = (1, 0, 0), v_2 = (0, 1, 0), v_3 = (0, 0, X), v_4 = (0, 0, Y), v_5 = (0, 0, Z)$. If $w_i \in \mathrm{H}^0(\mathbb{P}^2, Q)$ is the image of v_i under the isomorphism of (6) then $w_i, i = 1, \ldots, 5$ is a basis of $\mathrm{H}^0(\mathbb{P}^2, Q)$.

Lemma 4.1. *If E is vector bundle of rank 2 on $\mathbb{P}^2$ generated by three global sections then $c_1(E) = d$ and $c_2(E) = d^2$ for some integer $d \geq 0$.*

Proof: If a rank two vector bundle E is generated by three global sections then we get an exact sequence

$$0 \to \mathcal{O}_{\mathbb{P}^2}(-d) \to \mathcal{O}^3_{\mathbb{P}^2} \to E \to 0$$

of vector bundles on $\mathbb{P}^2$ and hence the required result. □

From Lemma(4.1) we see that the bundle Q on $\mathbb{P}^2$ that we have constructed above cannot be generated by 3 sections.

Example 1: Let Q be the vector bundle of rank two on $\mathbb{P}^2$ defined by the exact sequence (5). If $w_i, 1 \leq i \leq 5$ are the sections of Q defined above, then the set $S_1 = \{w_i; i = 1, \ldots, 4.\}$ is a generating set of sections of Q, i.e.,if $S = \{e_i; i = 1, \ldots, 4.\}$ is the standard basis of $\mathcal{O}^4_{\mathbb{P}^2}$ then the bundle map

$$\mathcal{O}^4_{\mathbb{P}^2} \to Q$$

obtained by sending e_i to w_i for $i = 1, \ldots, 4$ is surjective. Thus we get a morphism

$$\phi_{S_1} : \mathbb{P}^2 \to Gr(2, \mathbb{C}^4).$$

If $p : Gr(2, \mathbb{C}^4) \to \mathbb{P}^5$ is the Plucker imbedding, then the morphism

$$p \circ \phi_{S_1} : \mathbb{P}^2 \to \mathbb{P}^5$$

is given by $(x, y, z) \mapsto (z^2, -xy, -y^2, x^2, xy, 0)$. Let $z_i, (0 \leq i \leq 5)$ be the homogeneous coordinates of $\mathbb{P}^5$. Then the image of $\mathbb{P}^2$ under this morphism is a rank 3 quadric V in a linear $\mathbb{P}^3(\subset \mathbb{P}^5)$: In fact $V = Z(Z_5, Z_1 + Z_4, Z_1^2 + Z_2 Z_3))$. Let $p = (1, 0, 0, 0, 0, 0) \in V$ and $C = V \cap H$ where H is the hyperplane of $\mathbb{P}^5$ defined by Z_0. The morphism

$$\phi_{S_1}|_{\mathbb{P}^2 - \{(0,0,1)\}} =: \mathbb{P}^2 - \{(0, 0, 1)\} \to V - \{p\}$$

is a two sheeted ramified covering ramified precisely along C. Moreover $\phi_{S_1}((0, 0, 1)) = p$ and the differential map $d\phi_{S_1(0,0,1)}$ is zero. In this case we see that the image of $\mathbb{P}^2$ in $Gr(2, \mathbb{C}^4)$ is a singular surface with exactly one singularity.

Example 2: Let Q be the vector bundle of rank two on $\mathbb{P}^2$ defined by the exact sequence (5) and let $w_i, 1 \leq i \leq 5$ be the sections of Q defined above and let $u_1 = w_1, u_2 = w_2, u_3 = w_3, u_4 = w_4 - w_5$. The set $S_2 = \{u_i; i = 1, \ldots, 4\}$ is a generating set of sections of Q, i.e.,if $S = \{e_i; i = 1, \ldots, 4.\}$ is the standard basis of $\mathcal{O}^4_{\mathbb{P}^2}$ then the bundle map

$$\mathcal{O}^4_{\mathbb{P}^2} \to Q$$

obtained by sending e_i to u_i for $i = 1, \ldots, 4$, is surjectve. Thus we get a morphism

$$\phi_{S_2} : \mathbb{P}^2 \to Gr(2, \mathbb{C}^4).$$

If $p : Gr(2, \mathbb{C}^4) \to \mathbb{P}^5$ is the Plucker imbedding, then we see that the morphism

$$p \circ \phi_{S_2} : \mathbb{P}^2 \to \mathbb{P}^5$$

is given by

$$(x, y, z) \mapsto (z^2, -xy, -y^2 + yz, x^2, xy - xz, 0).$$

The image of $\mathbb{P}^2$ under this morphism is an intersection of two quadrics V in a linear $\mathbb{P}^4$: In fact $V = V(Z_5, Z_1Z_4 - Z_2Z_3, (Z_1 + Z_4)^2 - Z_0Z_3)$.

Example 3 : The vector space $\mathrm{H}^0(\mathbb{P}^2, \mathcal{O}^2_{\mathbb{P}^2} \oplus \mathcal{O}_{\mathbb{P}^2}(1))$ is equal to

$$\oplus_{i=1}^5 \mathbb{C}v_i,$$

where $v_1 = (1, 0, 0), v_2 = (0, 1, 0), v_3 = (0, 0, X), v_4 = (0, 0, Y), v_5 = (0, 0, Z)$. If $w_i \in \mathrm{H}^0(\mathbb{P}^2, Q)$ is the image of v_i under the isomorphism of (6) then $w_i, i = 1, \ldots, 5$ is a basis of $\mathrm{H}^0(\mathbb{P}^2, Q)$.

Let Q be the vector bundle of rank two on $\mathbb{P}^2$ defined by the exact sequence (5) and let $w_i, 1 \leq i \leq 5$ be the sections of Q defined above and let $u_1 = w_1, u_2 = w_2, u_3 = w_3 + dw_4, u_4 = aw_4 + w_5$, where a, d non zero complex numbers. The set $S_3 = \{u_i; i = 1, \ldots, 4\}$ is a generating set of sections of Q, i.e.,if $S = \{e_i; i = 1, \ldots, 4.\}$ is the standard basis of $\mathcal{O}^4_{\mathbb{P}^2}$ then the bundle map

$$\mathcal{O}^4_{\mathbb{P}^2} \to Q$$

obtained by sending e_i to u_i for $i = 1, \ldots, 4$, is surjectve. Thus we get a morphism

$$\phi_{S_3} : \mathbb{P}^2 \to Gr(2, \mathbb{C}^4).$$

If $p : Gr(2, \mathbb{C}^4) \to \mathbb{P}^5$ is the Plucker imbedding, then we see that the morphism

$$p \circ \phi_{S_3} : \mathbb{P}^2 \to \mathbb{P}^5$$

is given by

$$(x, y, z) \mapsto (z^2, -(x + dy)y, -(ay + z)y, (x + dy)x, (ay + z)x, 0).$$

The image of $\mathbb{P}^2$ under this morphism can be seen to be equal is equal to intersection of two independent singular quadrics in $\mathbb{P}^4$.

5. Main Theorem

Theorem 5.1. *Let $\phi : \mathbb{P}^2 \to Gr(2, \mathbb{C}^4)$ be a morphism. Assume that $c_1(Q) = 2$ and $c_2(Q) = 2$, where Q is the pull back by ϕ of the universal rank two quotient bundle on $Gr(2, \mathbb{C}^4)$. Then the image of $\mathbb{P}^2$ in $Gr(2, \mathbb{C}^4)$ is*

a) *either a complete intersection of two independent hyperplanes,*
b) *or is a complete intersection of a hyperplane and a quadric,*

Here a hyperplane(respectively, quadric) means a divisor in the class of the ample generator (respectively, twice the class of the ample generator) of the the Picard group of $Gr(2, \mathbb{C}^4)$. In case of a) the image is isomorphic to cone in $\mathbb{P}^3$ over a conic. In the case of b) the image is isomorphic to base locus of a pencil consists of singular quadrics in $\mathbb{P}^4$ of rank 3 and 4. More over the image surface is singular exactly along a line of $\mathbb{P}^4$ contained in the surface.

Proof: By our assumption, the vector bundle Q on $\mathbb{P}^2$ is generated by global sections. By Lemma 3.5, Q fits into an exact sequence (4). In the equation (4) the bundle map

$$0 \to \mathcal{O}_{\mathbb{P}^2}(-1) \to \mathcal{O}^2_{\mathbb{P}^2} \oplus \mathcal{O}_{\mathbb{P}^2}(1)$$

is given by $s = (A, B, Q)$, where $A, B \in \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^2}(1))$ and $Q \in \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^2}(2))$ with out common zeros in $\mathbb{P}^2$. Then the morphism ϕ is determined by four linearly independent global sections of the bundle $E = \mathcal{O}^2_{\mathbb{P}^2} \oplus \mathcal{O}_{\mathbb{P}^2}(1)$ whose images generate the bundle Q. If w_1, w_2, w_3, w_4 four linearly independent global sections of the bundle $E = \mathcal{O}^2_{\mathbb{P}^2} \oplus \mathcal{O}_{\mathbb{P}^2}(1)$ whose images generate the bundle Q and u_1, u_2, u_3, u_4 be any other basis of the vector space generated by w_1, w_2, w_3, w_4, then the morphism defined by w_1, w_2, w_3, w_4 and u_1, u_2, u_3, u_4 differ by an automorphism of $Gr(2, \mathbb{C}^4)$. Let $w_i = (a_i, b_i, g_i)$, $1 \leq i \leq 4$ be four global sections of $E = \mathcal{O}^2_{\mathbb{P}^2} \oplus \mathcal{O}_{\mathbb{P}^2}(1)$ such that their images in Q generate Q. Since Q is not direct sum of line bundles we see that sections of the bundle $\mathcal{O}^2_{\mathbb{P}^2}$ given by (a_i, b_i), $1 \leq i \leq 4$ generate $\mathcal{O}^2_{\mathbb{P}^2}$. Hence by taking suitable linear combinations we can assume that the four linearly independent global sections are of the form $w_1 = (1, 0, f_1), w_2 = (0, 1, f_2), w_3 = (0, 0, f_3), w_4 = (0, 0, f_4)$, where $f_3, f_4 \in \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^2}(1))$ are linearly independent. If $p : Gr(2, \mathbb{C}^4) \to \mathbb{P}^5$ is the Plucker imbedding, then we see that the morphism

$$p \circ \phi : \mathbb{P}^2 \to \mathbb{P}^5$$

is given by

$$(x; y; z) \mapsto$$

$$(D_0(x, y, z); D_1(x, y, z); D_2(x, y, z); D_3(x, y, z); D_4(x, y, z); 0),$$

where $D_0 = Q - f_1A, D_1 = -(f_3B), D_2 = -(f_4B), D_3 = -f_3A, D_4 = -f_4B$.

Case 1: The subspace $< f_3, f_4 >$ generated by f_3, f_4 is equal to $< A, B >$. In this case the four quadrics $-f_3B, -f_4B, f_3A, f_4A$ on $\mathbb{P}^2$ are linearly dependent, say $a(-f_3B)+b(-f_4B)+c(f_3A)+d(f_4A) = 0$, with $(a, b, c, d) \neq (0, 0, 0, 0)$. Then the image of $\mathbb{P}^2$ in $\mathbb{P}^5$ is an intersection of two independent hyperplanes namely, $Z_5 = 0$ and $aZ_1 + bZ_2 + cZ_3 + dZ_4 = 0$, with the Plucker embedding of the Grassmannnian given by $Z_0Z_5 + Z_1Z_4 - Z_2Z_3$, where $Z_0, \ldots, Z_5$ are the Plucker coordinate functions. Hence the image of $\mathbb{P}^2$ in $Gr(2, \mathbb{C}^4)$ is as stated in the case a) of the theorem.

Case 2: The subspace $< f_3, f_4 >$ generated by f_3, f_4 is not equal to $< A, B >$. In this case, if there is a non-trivial linear relation

$$a(Q - f_1B) + b(-f_3B) + c(-f_4B) + d(f_3A) + e(f_4A) = 0$$

exists, then again we see that the image of $\mathbb{P}^2$ in $\mathbb{P}^5$ under $p \circ \phi$ is an intersection of two independent hyperplanes namely, $Z_5 = 0$ and $aZ_0 + bZ_1 + cZ_2 + dZ_3 + eZ_4 = 0$ with the Plucker embedding of the Grassmannnian given by $Z_0Z_5 + Z_1Z_4 - Z_2Z_3$. Hence the image of $\mathbb{P}^2$ in $Gr(2, \mathbb{C}^4)$ is as stated in the case a) of the theorem.

Continuing the proof in the case 2, we can assume image is not as in a) in which case degree of the image of $\mathbb{P}^2$ in $\mathbb{P}^5$ under $p \circ \phi$ has to be four. As above set

$$D_1 = (Q - f_1B), D_2 = (-f_3B), D_3 = (-f_4B), D_4 = (f_3A), D_5 = (f_4A),$$

then we see that

$$D_2D_5 - D_3D_4 = 0. \tag{7}$$

This shows that image of $\mathbb{P}^2$ under $p \circ \phi$ is a degree four surface contained in intersection of the hyperplane $Z_5 = 0$ with the plucker embedding of the Grassmannian given by the equation $Z_0Z_5 + Z_1Z_4 - Z_2Z_3 = 0$, where $Z_i\,(0 \leq i \leq 5)$ are the homogeneous coordinate functions on $\mathbb{P}^5$. By Tango's result about the embeddings of $\mathbb{P}^2$ in Grassmannian (see, [11]) we conclude that the image of $\mathbb{P}^2$ under $p \circ \phi$ is singular surface of degree four in $\mathbb{P}^4$ and is given by base point free linear system of conics. Since the linear system is of dimension four we see that the image of $\mathbb{P}^2$ under $p \circ \phi$ is projection of the Veronese surface $V \subset \mathbb{P}^5$ from a point $p \in sec(V) \subset \mathbb{P}^5$ and $p \notin V$, where $sec(V)$ denotes the secant variety of V. Hence by Remark (2.1) it follows that the image of $\mathbb{P}^2$ in $\mathbb{P}^4$ given by $p \circ \phi$ is cut out by a linear pencil of quadrics in $\mathbb{P}^4$. Thus the image of $\mathbb{P}^2$ under ϕ in this case is a complete intersection of a hyperplane and a quadric as stated in b).

The last statement about the singularities can be easily checked. □

Acknowledgements: We thank referee for his valuable suggestions. Last named author would like to thank University of Lille-1 at Lille and University of Artois at Lens. He would also like to thank university of Paris 6 and IRSES-Moduli program.

References

[1] El Mazouni, A.; Laytimi, F.; Nagaraj, D. S. *Morphisms from $\mathbb{P}^2$ to $Gr(2, \mathbb{C}^4)$*. J. Ramanujan Math. Soc. 26 (2011), no. 3, 321–332.

[2] Mauro C. Beltrametti.; Ettore Carletti.; Dionisio Gallarati.; Giacomo Monti Bragadin. *Lectures on Curves, Surfaces and Projective Varieties Translated from the Italian by Francis Sullivan.* European Mathematical Society First corrected reprint, June 2012

[3] Sierra, Jos Carlos; Ugaglia, *Luca On double Veronese embeddings in the Grassmannian G(1,N).* Math. Nachr. 279 (2006), no. 7, 798–804.

[4] Sierra, Jos Carlos; Ugaglia, Luca *On globally generated vector bundles on projective spaces.* J. Pure Appl. Algebra 213 (2009), no. 11, 2141–2146.

[5] PH. Ellia. *Chern Classes of rank two globally generated vector bundles on $\mathbb{P}^2$*. arXiv:1111.5718, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. 24 (2013), no. 2, 147–163.

[6] W. Fulton and J. Harris. *Representation Theorey* GTM 129,Springer–Verlag.

[7] R. Hartshorne. *Stable Vector Bundle of Rank* 2 *on $\mathbb{P}^3$*. Math. Ann. 238, 229–280 (1978)

[8] R. Hartshorne. *Stable Reflexive Sheaves*, Math. Ann. 254, 121–176 (1980)

[9] R. Hartshorne. *Varieties of small codimension in projective space*, Bulletin of the American Mathematical Society 80: 1017–1032

[10] J. Harris. *Algebraic Geometry* GTM 133,Springer- -Verlag.

[11] Hiroshi Tango: *On (n-1)-dimensional projective spaces contained in the Grassmann variety $Gr(n, 1)$*. J. Math. Kyoto Univ. 14-3 (1974) 415-460.

Laboratoire de Mathématiques de Lens EA 2462 Faculté des Sciences Jean Perrin Rue Jean Souvraz, SP18 F-62307 LENS Cedex France

E-mail address: mazouni@euler.univ-artois.fr

F. L.: Mathématiques - bât. M2, Université Lille 1, F-59655 Villeneuve d'Ascq Cedex, France

E-mail address: fatima.laytimi@math.univ-lille1.fr

Institute of Mathematical Sciences C.I.T. campus, Taramani, chennai 600113,India

E-mail address: dsn@imsc.res.in